Web Images Videos Maps News Shopping Gmail more ▼ professorwhaley@gmail.com | Scholar Preferences | My Account | Sign out Google scholar weighted regression, disease risk model Search Advanced Service Search Scholar Articles and patents -- 2001 include citations R Create email alert Results 1 - 10 of about 18,000, (0.04 sec) Does the relation of blood pressure to coronary heart disease risk change with aging?: The ahajournals.org [HTML] Framingham Heart Study \$5. Frankin, MG Carson, SA Nhan, ND Wong, EP ... - Circulation, 2001. Am Heart Assoc ... Each examination included an extensive cardiovascular disease history and physical examination. 12-lead ... reliher SBP (P-0.07) nor DBP (P-0.74) precided CHD risk, possibly because ... The estimated slope per decade of age from the weighted regression procedure was 0.29 ... Cherg tw 690 - Paletier una les - St. Oines - An 5 venerors. Moderate alcohol intake and lower risk of coronary heart disease; meta-analysis of effects on lipids bmi.com [HTNK] and haemostatic factors
EB Rimm, P Williams, K Foster, M Criqui, MJ ... - British Medical ..., 1999 - projection 1, factor VII, and von Willebrand factor (see table B on website) factors that have been associated with risk of coronary heart disease. 26-28 Weighted regression analyses for each of these haemostatic factors was suggestive of a more thrombolytic profile, but none was ... Checky 886 - Palanet strates - 61, Oirech - Alt 9 ventrors. Insulin and risk of cardiovascular disease: a meta-analysis
JB Rugo, WJJ Asendein: JB Delker PJ Kosense, ... - Cito-lation 199- Am Heart Assor,
... The weight of each study was calculated inversely to the variance estimate of the ... in insulin level
in a Cox proportional hazards, logistic tragression, or Poisson tragression model. 12 The ebajournals.org [HTML] reexpressed RR of cardiovascular disease per specified uniform difference in insulin level is ... Cried by 324 - Reisted articles - Bt. Einect - At 10 versions nih.gov (PDF) Airborne particles are a risk factor for hospital admissions for heart and lung disease. Full View Ched by 198 - Related anivies - BL Dys.c. - All 29 versions Binomial regression in GLIM, estimating risk ratios and risk differences S WACHOLDER: American Journal of Epidemiology, 1966 - Oxford Univ Press the logarithm of the "health ratio" or the ratio of the probabilities of no disease at values of X ... Generally, when A and B are independent risk factors, absence of interaction on one scale implies ... If the TT.'S were known, a weighted regression could be used, with each obser- vation ... Cated by 250 - Related articles - At 6 versions [PDF] Depressed affect, hopelessness, and the risk of ischemic heart disease in a cohort of US istor.org (PDF) R Anda O Williamson, O Jones, C Macera, E Eaker, A . - Epidemology, 1993 - JSTOP ... fitting a weighted (using the inverse of the standard error as the weight) regression line through ... 40%, respectively, for each increase in level of hope, lessness (the regression analysis for ... early mortally that may have occurred because of unidentified or occult disease at baseline. Cited by 556 - Related articles - Rt. Ohect - At 6 versions Effect of prednisone and hydroxychloroquine on coronary artery disease risk factors in systemic lupus erythematosus: a longitudinal data analysis
M Petri, C Liskatta, L Magder, D Goldman - The American lournal of ..., 1994 - Bisever .. why treated hypertension remains a risk factor in our model for coronary artery disease [18] ... that an increase in prednisone dose frequently, almost inexorably, leads to an increase in weight. Regression analysis of longitudinal data may not be the best way to model weight. ... Cited by 160 - Related articles - At 3 yerskins Spatial variation in risk of disease; a nonparametric binary regression approach ... Fit a weighted additive model of the form Z uH g x, using kernel regression with weights wi as follows. (i) Put siz  $\hat{A}$  uH i  $\hat{e}$  and perform a weighted kernel regression of  $\hat{s}$  to  $\hat{n}$  x with weights wi eg x  $\hat{n}$  1 in w Kh x  $\hat{A}$  x  $\hat{n}$  0  $\hat{n}$  1 if  $\hat{n}$  k  $\hat{n}$  5 x  $\hat{n}$  6 x  $\hat{n}$  5 x  $\hat{n}$  5 x  $\hat{n}$  6 x  $\hat{n}$  7 x  $\hat{n}$  6 x  $\hat{n}$  6 x  $\hat{n}$  7 x  $\hat{n}$  7 x  $\hat{n}$  7 x  $\hat{n}$  8 x  $\hat{n}$  7 x  $\hat{n}$  8 x  $\hat{n$ Gred by 118 - Resided articles - Bi, Deed - All 6 versions Blood pressure, stroke, and coronary heart disease\* 1; Part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression ... .. vertical line represents a 29 difference in risk, which was the weighted mean difference in ... of multiple regression analysis; and regression to the mean (and regression from the ... intake associated with elevated blood pressure: the Kuopio ischaemic heart disease risk factor study ... Coad by 2372 - Resided enjoyee - All 5 versions High risk of thrombosis in patients homozygous for factor V Leiden (activated protein C resistance) hematology/ib/rary.org [PDF] FR Resentiaal, T Koster, JP ....., 1995 - bloodysumal.nemetslogy/ibrary.org .. Angina pectoris, previous myocardial infarction, stroke, or peripheral arterial disease. ... Therefore, as an overall figure, the incidence figures that were recalculated from the weighted regression model seem the best estimate of the risk, which becomes over 2% per year in the ... Cond by 757 - Resided artists - Bt. Direct - At 12 persions

Create email alert

Gonooooooong ic ➤
Result Page: 12345678910 Next

weighted regression, disease risk m Search

Go to Google Home - About Google - About Google Scholair

©2010 Google

/PW/